

Публикации по теме

Kuznetsova I. M., Biktashev A.G., Malova L.N., Bushmarina N.A., Uversky V.N., and Turoverov K.K. 2000. Understanding the contribution of individual tryptophan residues to intrinsic lysozyme fluorescence. *Protein and Peptide Letters*. 7, 6: 411–420.

[Kuznetsova I.M., Yakusheva T.A., Turoverov K.K.](#) 1999. Contribution of separate tryptophan residues to intrinsic fluorescence of actin. Analysis 3D structure. *FEBS Lett.* 452: 205–210.

[Кузнецова И.М., Туроверов К.К.](#) 1998. Что определяет характеристики собственной УФ-флуоресценции белков? Анализ свойств микроокружения и особенностей локализации их триптофановых остатков. *Цитология*. 40 (8/9): 747–762.

[Turoverov K.K., and Kuznetsova I.M.](#) 1986. What causes the depolarization of trypsin and trypsinogen fluorescence. Intramolecular mobility or non-radiative energy transfer? *Biophys.Chem.* 25: 315–323.

[Turoverov K. K., Kuznetsova I. M., and Zaitsev V. N.](#) 1985. The environment of the tryptophan residue in *Pseudomonas aeruginosa* azurin and its fluorescence properties. *Biophys. Chem.* 23: 79–89.