

PUBLICATIONS

Povarova O.I., Sulatskaya A.I., Kuznetsova I.M., Turoverov K.K. 2012. The interaction of the hydrophobic dye ANS with inactivated actin. FEBS Journal 279 (Suppl. 1). 279: 415.

[Povarova O.I., Kuznetsova I.M. and Turoverov K.K.](#) 2010. Differences in the pathways of proteins unfolding induced by urea and guanidine hydrochloride: molten globule state and aggregates. PLoS ONE 5(11): e15035. doi: 10.1371/journal.pone.0015035.

[Povarova O.I., Kuznetsova I.M., Turoverov K.K.](#) 2007. Different disturbances – one pathway of protein unfolding. Actin folding-unfolding and misfolding. Cell.Biol.Int. 31, 405-412. doi: 10.1016/j.cellbi.2007.01.025.

[Mazhul' V.M., Zaitseva E.M., Shavlovsky M.M., Stepanenko Olesia V., Kuznetsova I.M., and Turoverov K.K.](#) 2003. Monitoring of Actin Unfolding by Room Temperature Tryptophan Phosphorescence. Biochemistry. 42, 13551-13557. doi: 10.1021/bi0350295

Turoverov K.K. and Kuznetsova I.M.. 2003 Intrinsic fluorescence of actin. J. Fluorescence. 13, 41-57. doi: 10.1023/A:1022366816812.

[Kuznetsova I.M., Stepanenko Olga V., Stepanenko Olesia V., Povarova O.I., Biktashev A.G., Verkhusha V.V., Shavlovsky M.M., and Turoverov K.K.](#) 2002. The place of inactivated actin and its kinetic predecessor in actin folding-unfolding. Biochemistry 41, 13127-13132.

[Turoverov K.K., Verkhusha V.V., Shavlovsky M.M., Biktashev A.G., Povarova O.I., and Kuznetsova I.M.](#) 2002. Kinetics of actin unfolding induced by guanidine hydrochloride. Biochemistry 41, 1014-1019.

[Kuznetsova I.M., Biktashev A.G., Khaitlina S.Yu., Vassilenko K.S., Turoverov K.K. and Uversky V.N.](#) 1999. Effect of self-association on the structural organization of partially folded proteins: inactivated actin. Biophys.J. 77, 5:2788-2800. doi: 10.1016/S0006-3495(99)77111-0.

[Turoverov K.K., Biktashev A.G., Khaitlina S.Yu., and Kuznetsova I.M.](#) 1999. The structure and dynamics of partially-folded actin. Biochemistry. 38:6261-6269. doi: 10.1021/bi9900976

Kuznetsova I.M., Turoverov K.K. and Uversky V.N. 1999. Inactivated actin, and aggregate comprised of partially-folded monomers, has a overall native-like packing density. Protein and Peptide Letters. 6:173-178.

Turoverov K.K., Kuznetsova I.M., Khaitlina S.Yu., and Uversky V.N. 1999. Unusual combination of the distorted structure and frozen mobility in inactivated actin molecule. Protein and Peptide Letters. 6:73-78.

[Kuznetsova I.M., Khaitlina S.Y., Konditerov S.N., Surin A.M., and Turoverov K.K.](#) 1988. Changes of the structure and intramolecular mobility in the course of actin denaturation. Biophys.Chem. 32:73-78.

[Turoverov K.K., Haitlina S.Y., and Pinaev G.P.](#) 1976. Ultra-violet fluorescence of actin. Determination of native actin content in actin preparations. FEBS Letters. 62: 4-6.