

Публикации по теме

Lavysh A. V., Maskevich A. A., Lugovskii A. A., Voropay E. S., Sulatskaya A. I., Kuznetsova I. M., Turoverov K. K. Formation of trans-2-[4-(Dimethylamino) Styryl]-3-Ethyl-1, 3-Benzothiazolium Perchlorate Dimers in the Presence of Sodium Polystyrene Sulfonate. 2017. Journal of Applied Spectroscopy. 83(6): 917-923. doi: 10.1007/s10812-017-0384-8.

[Rodina N.P., Sulatsky M.I., Sulatskaya A.I., Kuznetsova I.M., Uversky V.N., Turoverov K.K.](#) 2017. Photophysical properties of fluorescent probe thioflavin T in crowded milieu. Journal of Spectroscopy. Volume 2017, Article ID 2365746, 10 pages. doi: 10.1155/2017/2365746.

[Sulatskaya A.I., Rodina N.P., Povarova O.I., Kuznetsova I.M., Turoverov K.K.](#) 2017 Different conditions of fibrillogenesis cause polymorphism of lysozyme amyloid fibrils. Journal of Molecular Structure. 1140:52-58. doi: [10.1016/j.molstruc.2016.10.037](#).

[Lavysh A. V., Lugovskii A. A., Voropay E. S., Sulatskaya A. I., Kuznetsova I. M., Turoverov K. K., Maskevich A. A.](#) 2016. Aggregation of thioflavin T and its new derivative in the presence of anionic polyelectrolyte. Biointerface Research in Applied Chemistry. 6 (5): 1525-1530.

[Kuznetsova I.M., Sulatskaya A.I., Maskevich A.A.; Uversky V.N.; Turoverov K.K.](#) 2016. The high fluorescence anisotropy of thioflavin T in aqueous solution results from its molecular rotor nature. Analytical Chemistry. 88(1): 718-724. doi: 10.1021/acs.analchem.5b02747.

[Sulatskaya A.I., Kuznetsova I.M., Belousov M.V., Bondarev S.A., Zhouravleva G.A., Turoverov K.K.](#) 2016. Stoichiometry and Affinity of Thioflavin T Binding to Sup35p Amyloid Fibrils. PLoS One. 11(5): e0156314. doi: 10.1371/journal.pone.0156314.

[Sulatskaya A.I., Kuznetsova I.M., Turoverov K.K.](#) 2016. Photophysical properties of thioflavin T. Does it form excimers when integrated into amyloid fibrils? Biophys. J. 110 (3): 218a. doi: [10.1016/j.bpj.2015.11.1210](#)

Maskevich A.A., Lavysh A.V., Kuznetsova I.M., Sulatskaya A.I., Turoverov K.K. 2015. Spectral Manifestations of Thioflavin T Aggregation. Journal of Applied Spectroscopy. 82 (1): 33-39. doi: 10.1007/s10812-015-0060-9.

[Fonin A.V., Sulatskaya A.I., Kuznetsova I.M., Turoverov K.K.](#) 2014. Fluorescence of dyes in solutions with high absorbance. Inner filter effect correction. Plos One. 9(7): e103878. doi: 10.1371/journal.pone.0103878.

Lavysh A.V., Sulatskaya A.I., Lugovskii A.A., Voropay E. S., Kuznetsova I.M., Turoverov K.K., Maskevich A.A. 2014. Photophysical properties of trans-2-[4-(dimethylamino)styryl]-3-ethyl-1,3-benzothiazolium perchlorate, a new structural analog of thioflavin T. J. Appl. Spectr. 81(2): 205-213. doi: 10.1007/s10812-014-9911-z.

[Сулацкая А. И., Волова Е. А., Комиссарчик Я.Ю., Снигиревская Е.С., Маскевич А.А., Дробченко Е.А., Кузнецова И.М., Тuroverov K.K.](#) 2013. Исследование кинетики образования амилоидных фибрилл на основе инсулина. Цитология 55(11): 807-812.

[Kuznetsova Irina M., Sulatskaya Anna I., Povarova Olga I., Turoverov Konstantin K.](#) 2012. Reevaluation of ANS binding to Human and Bovine Serum Albumins. Key Role of Equilibrium Microdialysis in Ligand - Receptor Binding Characterization. Plos One. 7(7): e40845. doi: 10.1371/journal.pone.0040845.

[Kuznetsova I.M., Sulatskaya A. I., Uversky V.N., Turoverov K.K.](#) 2012. A new trend in the experimental methodology for the analysis of the Thioflavin T binding to amyloid fibrils. Molecular Neurobiology. 45: 488-498. doi: 10.1007/s12035-012-8272-y.

[Sulatskaya A.I., Kuznetsova I.M., Turoverov K.K.](#) 2012. Interaction of thioflavin T with amyloid fibrils: fluorescence quantum yield of bound dye. The Journal of physical chemistry B. 116(8): 2538-2544. doi: 10.1021/jp2083055.

[Kuznetsova I.M., Sulatskaya A.I., Uversky V.N., Turoverov K.K.](#) 2012. Analyzing Thioflavin T binding to amyloid fibrils by an equilibrium microdialysis-based technique. Plos One. 7(2): e30724. doi: 10.1371/journal.pone.0030724.

[Sulatskaya A.I., Kuznetsova I.M., Turoverov K.K.](#) 2011. Interaction of thioflavin T with amyloid fibrils: stoichiometry and affinity of dye binding, absorption spectra of bound dye. J. Phys. Chem. B. 115 (39): 11519-11524. doi: 10.1021/jp207118x.

[Stsiapura V.I., Maskevich A.A., Kuzmitsky V.A., Uversky V.N., Kuznetsova I.M., Turoverov K.K.](#) 2008. Thioflavin T as a molecular rotor: fluorescent properties of thioflavin T in solvents with different viscosity. J.Phys.Chem. B . 112(49) 15893-15902. doi: 10.1021/jp805822c.

[Maskevich A.A., Stsiapura V.I., Kuzmitsky V.A., Kuznetsova I.M., Povarova O.I., Uversky V.N. and Turoverov K.K.](#) 2007. Spectral properties of thioflavin T in solvents with different dielectric properties and in fibril-incorporated form. J. Proteome Res. 6(4), 1392-1401. doi: [10.1021/pr0605567](#).