

PUBLICATIONS

- Stepanenko Olga V., Roginskii D.O., Stepanenko Olesya V., Kuznetsova I.M., Uversky V.N., Turoverov K.K. 2016. Structure and stability of recombinant bovine odorant-binding protein: I. Design and analysis of monomeric mutants. PeerJ. 4: e1933. doi: 10.7717/peerj.1933.
- Stepanenko Olga V., Roginskii D.O., Stepanenko Olesya V., Kuznetsova I.M., Uversky V.N., Turoverov K.K. 2016. Structure and stability of recombinant bovine odorant-binding protein: II. Unfolding of the monomeric forms. PeerJ. 4: e1574. doi: 10.7717/peerj.1574.
- Stepanenko Olga V., Fonin A.V., Stepanenko Olesya V., Staiano M., D'Auria S., Kuznetsova I.M., Turoverov K.K. 2015. Tryptophan residue of the D-galactose/D-glucose-binding protein from E. Coli localized in its active center does not contribute to the change in intrinsic fluorescence upon glucose binding. J. Fluorescence 25(1):87-94. doi: 10.1007/s10895-014-1483-z.
- Fonin A.V., Povarova O.I., Staiano M., D'Auria S., Turoverov K.K., Kuznetsova I.M. 2014. Trehalose/maltose-binding protein as a sensitive element of glucose biosensor. Optical Materials 36(10): 1676 – 1679. [10.1016/j.optmat.2014.01.001](https://doi.org/10.1016/j.optmat.2014.01.001).
- Stepanenko Olga V., Stepanenko Olesya V., Staiano M., Kuznetsova I.M., Turoverov K.K., D'Auria S. 2014. The Quaternary Structure of the Recombinant Bovine Odorant-Binding Protein is Modulated by Chemical Denaturants. PlosOne. 9(1): e85169. doi: 10.1371/journal.pone.0085169.
- Fonin A.V., Stepanenko O.V., Povarova O.I., Volova C.A., Philippova E.M., Bublikov G.S., Kuznetsova I.M., Demchenko A.P., Turoverov K.K. 2014. Spectral characteristics of the mutant form GGBP/H152C of D-glucose/D-galactose-binding protein labeled with fluorescent dye BADAN: influence of external factors. PeerJ. 2: e275. doi: 10.7717/peerj.275.
- Stepanenko Olga V., Stepanenko Olesya V., Povarova O.I., Fonin A.V., Kuznetsova I.M., Turoverov K.K., Staiano M., Varriale A., D'Auria S. 2011. New insight into protein-ligand interactions. The case of the D-galactose/D-glucose-binding protein from Escherichia coli. J. Phys. Chem. B. 115 (12): 2768-73. doi: 10.1021/jp1095486.
- Stepanenko Olga V., Fonin A.V., Stepanenko Olesya V., Morozova K.S., Verkhusha V.V., Kuznetsova I.M., Turoverov K.K., Staiano M., D'Auria S. 2011. New insight in protein-ligand interactions. 2. Stability and properties of two mutant forms of the D-galactose/D-glucose-binding protein from E. coli. J. Phys. Chem. B. 115 (29): 9022-32. doi: 10.1021/jp204555h.
- Stepanenko Olesya V., Marabotti A., Kuznetsova I. M., Turoverov K.K., Fini C., Varriale A., Staiano M., Rossi M., and D'Auria S. 2008. Hydrophobic amino acid residues buried in the matrix of the porcine odorant-binding protein play an important role in protein stability and dynamics. Proteins. 71(1): 35-44. doi: 10.1002/prot.21658.
- Staiano M., D'Auria S., Varriale A., Rossi M., Marabotti A., Fini C., Stepanenko Olesya V., Kuznetsova I. M., and Turoverov K.K. 2007. Stability and Dynamics of the Porcine Odorant-Binding Protein. Biochemistry 46(39):11120-7. doi: 10.1021/bi7008129.
- Kuznetsova I.M., Povarova O., Stepanenko Olesya, Turoverov K.K., Crescenzo R., Varriale A., Staiano M., and D'Auria S. 2007. New perspectives in protein-based biosensors: The glucokinase from *B. stearothermophilus* and the odorant-binding protein from *C. familiaris* as probes for non-consuming analyte sensors. Proc. SPIE, 6733, 673318, 1-6. doi: 10.1117/12.753249.

Kuznetsova I.M., Stepanenko Olga V., Turoverov K.K., Scognamiglio V., Staiano M., Rossi M., D'Auria S. 2005. Fluorescence properties of glutamine-binding protein from Escherichia coli and its complex with glutamine. J. Proteome Res. 4, 417-423. doi: 10.1021/pr0498077

Staiano M., Scognamiglio V., Rossi M., D'Auria S., Stepanenko Olga V., Kuznetsova I.M., Turoverov K.K. 2005. Unfolding and refolding of the glutamine-binding protein from Escherichia coli and its complex with glutamine induced by guanidine hydrochloride. Biochemistry. 44(15): 5625-5633. doi: 10.1021/bi0478300.