

Ion Channels in Cell Proliferation and Apoptotic Cell Death

F. Lang¹, M. Föller¹, K.S. Lang¹, P.A. Lang¹, M. Ritter², E. Gulbins¹, A. Vereninov³, S.M. Huber¹

¹Department of Physiology, University of Tübingen, Germany

²Department of Physiology, University of Salzburg, Austria

³Institute of Cytology, Russian Academy of Sciences, St. Petersburg, Russia

Received: 9 September 2005

Abstract. Cell proliferation and apoptosis are paralleled by altered regulation of ion channels that play an active part in the signaling of those fundamental cellular mechanisms. Cell proliferation must - at some time point - increase cell volume and apoptosis is typically paralleled by cell shrinkage. Cell volume changes require the participation of ion transport across the cell membrane, including appropriate activity of Cl^- and K^+ channels. Besides regulating cytosolic Cl^- activity, osmolyte flux and, thus, cell volume, most Cl^- channels allow HCO_3^- exit and cytosolic acidification, which inhibits cell proliferation and favors apoptosis. K^+ exit through K^+ channels may decrease intracellular K^+ concentration, which in turn favors apoptotic cell death. K^+ channel activity further maintains the cell membrane potential, a critical determinant of Ca^{2+} entry through Ca^{2+} channels. Cytosolic Ca^{2+} may trigger mechanisms required for cell proliferation and stimulate enzymes executing apoptosis. The switch between cell proliferation and apoptosis apparently depends on the magnitude and temporal organization of Ca^{2+} entry and on the functional state of the cell. Due to complex interaction with other signaling pathways, a given ion channel may play a dual role in both cell proliferation and apoptosis. Thus, specific ion channel blockers may abrogate both fundamental cellular mechanisms, depending on cell type, regulatory environment and condition of the cell. Clearly, considerable further experimental effort is required to fully understand the complex interplay between ion channels, cell proliferation and apoptosis.

Key words: CD95/Fas — Scramblase — PGE2 — Cell volume — Lymphocytes — Erythrocytes

Introduction

Cell homeostasis requires a delicate balance between formation of new cells by cell proliferation and their elimination by apoptosis. Apoptosis eliminates abundant and potentially harmful cells (Green & Reed, 1998). The maintenance of an adequate cell number requires the replacement of apoptotic cells or formation of additional cells by cell proliferation.

Cell proliferation is stimulated by growth factors (Bikfalvi et al., 1998; Adams et al., 2004; Tallquist & Kazlauskas, 2004), apoptosis is triggered by stimulation of respective receptors, such as CD95 (Lang et al., 1998b, 1999; Gulbins et al., 2000; Fillon et al., 2002), somatostatin receptor (Teijeiro et al., 2002) or TNF α receptor (Lang et al., 2002a), by cell density (Long et al., 2003), lack of growth factors (Sturm et al., 2004) thyroid hormones (Alia et al., 2005), or adhesion (Davies, 2003; Walsh et al., 2003), by choline deficiency (Albright et al., 2005), DNA damage (Kohn & Pommier, 2005), or by exposing cells to genotoxic or other stressors such as radiation (Rosette & Karin, 1996), chemotherapeutics (Carriers et al., 2002; Wieder et al., 2001), oxidants (Rosette & Karin, 1996), inhibition of glutaminase (Rotoli et al., 2005), energy depletion (Pozzi et al., 2002) or osmotic shock (Bortner & Cidlowski, 1998, 1999; Lang et al., 1998a, 2000b; Maeno et al., 2000; Michea et al., 2000; Rosette & Karin, 1996).

If cell proliferation is to generate cells similar to the parent cells, it obviously requires the duplication of all cell components, such as DNA, cytoskeleton, mitochondria, etc. To generate cells of similar size as the parent cells, cell proliferation must at some point lead to cell volume increase (Lang et al., 1998a).

Hallmarks of apoptosis include nuclear condensation, DNA fragmentation, mitochondrial depolarization, cell shrinkage and breakdown of

phosphatidylserine asymmetry of the plasma membrane (Green & Reed, 1998). The exposure of phosphatidylserine at the cell surface results from the activation of a scramblase (Zhou et al., 2002), which is activated by increase of cytosolic Ca^{2+} activity (Woon et al., 1999; Dekkers et al., 2002). As macrophages are equipped with receptors specific for phosphatidylserine (Fadok et al., 2000; Henson et al., 2001), cells or cellular fragments exposing phosphatidylserine at their surface will be rapidly recognized, engulfed and degraded (Boas et al., 1998). Cell shrinkage facilitates the engulfment of the dying cells by phagocytes (Boas et al., 1998). Thus, apoptosis allows the elimination of the cells without release of intracellular proteins, which would otherwise cause inflammation (Gulbins et al., 2000).

Programmed cell death is not limited to nucleated cells but could similarly affect erythrocytes (Barvitenko et al., 2005; Bosman & Willekens, 2005; Rice & Alfrey, 2005). Recently, the term “eryptosis” has been coined (Lang et al., 2005a) to describe apoptosis-like death of mature erythrocytes, which is characterized by cell shrinkage and breakdown of phosphatidylserine asymmetry, both typical features of apoptosis in nucleated cells (Lang et al., 2003a, b, c, e).

Both cell proliferation and apoptosis involve at some point activation of Cl^- channels, K^+ channels and Ca^{2+} channels. As the respective channel inhibitors have been reported to interfere with both cell proliferation and apoptosis, the channels appear to play an active role in the machinery leading to duplication or death of a given cell.

The Role of Ca^{2+} and Nonselective Cation Channels in Cell Proliferation and Apoptosis

Overwhelming evidence points to a critical role of cytosolic Ca^{2+} activity in the regulation of cell proliferation (Whitfield et al., 1995; Parekh & Penner, 1997; Berridge et al., 1998, 2000, 2003; Santella 1998; Santella et al., 1998). Growth factors are known to stimulate Ca^{2+} release-activated calcium channels I_{CRAC} (Qian & Weiss, 1997), which has in turn been shown to trigger Ca^{2+} entry into and subsequent Ca^{2+} oscillations in proliferating cells. The Ca^{2+} oscillations trigger a wide variety of cellular functions (Berridge et al., 1998, 2000, 2003; Parekh & Penner, 1997), including the depolymerization of the actin filaments (Lang et al., 1992, 2000c; Dartsch et al., 1995; Ritter et al., 1997). The depolymerization of the actin filaments leads to disinhibition of the Na^+/H^+ exchanger and/or the Na^+ , K^+ , 2Cl^- cotransporter (Fig. 1) and thus leads to increase of cell volume (Lang et al., 1998a). Activation of I_{CRAC} , Ca^{2+} oscillations and depolymerization of the actin filament network all have been shown to be prerequisites of cell proliferation (Dartsch et al., 1995; Lang et al., 1992, 2000c; Ritter et al., 1997).

Conversely, the lymphocyte apoptosis following CD95 receptor triggering is paralleled by inhibition of I_{CRAC} (Lepple-Wienhues et al., 1999; Dangel et al., 2005) (Fig. 2). Inhibition of I_{CRAC} presumably serves to abrogate activation and proliferation of lymphocytes and does not necessarily foster apoptotic cell death. Whether or not cytosolic Ca^{2+} activity increases at a later stage following CD95 triggering, remains uncertain. In any case, increase of cytosolic Ca^{2+} is not an early event in CD95-induced cell death.

On the other hand, sustained increase of cytosolic Ca^{2+} activity has been shown to trigger apoptosis in a variety of nucleated cells (Parekh & Penner, 1997; Green & Reed, 1998; Berridge et al., 2000; Spassova et al., 2004; Liu et al., 2005; Parekh & Putney, Jr., 2005). Moreover, as illustrated in Fig. 3, Ca^{2+} -permeable cation channels play a decisive role in apoptosis-like death of erythrocytes (eryptosis) (Brand et al., 2003; Lang et al., 2002b; Lang et al., 2003b).

Apoptosis-like erythrocyte death could be triggered by exposure to the Ca^{2+} ionophore ionomycin (Berg et al., 2001; Bratosin et al., 2001; Daugas et al., 2001; Lang et al., 2002b, 2003b) and blunted in the nominal absence of Ca^{2+} (Lang et al., 2003b). The Ca^{2+} -permeable cation channels could be activated by exposure of erythrocytes to osmotic shock (Huber et al., 2001), oxidative stress (Duranton et al., 2002), energy depletion (Lang et al., 2003b) and infection with the malaria pathogen *Plasmodium falciparum* (Brand et al., 2003; Duranton et al., 2003; Lang et al., 2004a). Energy depletion presumably impairs the replenishment of GSH and thus weakens the antioxidative defense of the erythrocytes (Mavelli et al., 1984; Bilmen et al., 2001). The cation channels are further activated by removal of intracellular and extracellular Cl^- (Huber et al., 2001; Duranton et al., 2002). The suicidal cation channels are probably identical to the Na^+ and K^+ permeability activated by incubation of human erythrocytes in low ionic strength (LaCelle & Rothsteto, 1966; Jones & Knauf, 1985; Bernhardt et al., 1991). Similar if not identical nonselective cation channels are activated by depolarization (Christophersen & Bennekou, 1991; Bennekou, 1993; Kaestner et al., 1999).

Elevated cytosolic Ca^{2+} concentrations then stimulate the erythrocyte scramblase (Zhou et al., 2002), thus leading to the breakdown of phosphatidylserine asymmetry (Lang et al., 2003b). The phosphatidylserine exposure following osmotic shock is blunted by amiloride (Lang et al., 2003b) and ethylisopropylamiloride (EIPA) (Lang et al., 2003a), both putative inhibitors of the cation channel (Lang et al., 2003b, c). The suicidal erythrocyte cation channels are activated by prostaglandin E_2 (PGE_2), which is released upon osmotic shock (Lang et al., 2005b).

Cell volume-sensitive cation channels are expressed in a wide variety of nucleated cells, such as airway epithelia cells (Chan et al., 1992), mast cells




Fig. 1. Involvement of channels in the activation of cell proliferation by ras oncogene. In ras oncogene-expressing cells the mitogen bradykinin triggers the formation of inositol-(1,4,5) trisphosphate (IP₃) with subsequent stimulation of intracellular Ca²⁺ release. The subsequent activation of the Ca²⁺ release-activated Ca²⁺ channel *I*_{CRAC} leads to Ca²⁺ entry. Ca²⁺ activates Ca²⁺-sensitive K⁺ channels leading to K⁺ exit, hyperpolarization, and Cl⁻ exit through Cl⁻ channels. The subsequent cell shrinkage is required for the initial triggering of Ca²⁺ and cell membrane potential oscillations due to repetitive Ca²⁺ and K⁺ channel activation. The oscillations of Ca²⁺ lead to depolymerization of the cell actin filaments, which disinhibits the Na⁺/H⁺ exchanger and the Na⁺, K⁺, 2Cl⁻ cotransporter. Electrolyte accumulation by these carriers eventually leads to cell swelling.

(Cabado et al., 1994), macrophages (Gamper et al., 2000), vascular smooth muscle, colon carcinoma and neuroblastoma cells (Koch & Korbmaier, 1999), cortical collecting duct cells (Volk et al., 1995), and hepatocytes (Wehner et al., 1995, 2000). Cation channels activated by Cl⁻ removal were identified in salivary and lung epithelial cells (Marunaka et al., 1994; Tohda et al., 1994; Dinudom et al., 1995). It has been shown that Cl⁻ influences the channels via a pertussis toxin-sensitive G-protein (Dinudom et al., 1995). At present, we do not know the molecular identity of those channels nor do we know whether or not they participate in Ca²⁺ entry and apoptosis.

The Role of K⁺ Channels in Cell Proliferation and Apoptosis

A variety of K⁺ channels have been implicated in the regulation of cell proliferation (Patel & Lazdunski,

2004; Wang, 2004). Growth factors have been shown to activate K⁺ channels (O'Lague et al., 1985; Enomoto et al., 1986; Lang et al., 1991; Sanders et al., 1996; Wiecha et al., 1998; Liu et al., 2001; Faehling et al., 2001;), and enhanced K⁺ channel activity has been observed in a wide variety of tumor cells (DeCoursey et al., 1984; Nilius & Wohlrab, 1992; Pappone & Ortiz-Miranda, 1993; Strobl et al., 1995; Mauro et al., 1997; Skryma et al., 1997; Pappas & Ritchie, 1998; Zhou et al., 2003; Patel & Lazdunski, 2004; Wang, 2004). As illustrated in Fig. 1, repetitive activation of Ca²⁺-sensitive K⁺ channels by oscillating cytosolic Ca²⁺ activity leads to oscillations of cell membrane potential in ras oncogene-expressing cells (Lang et al., 1991). Several K⁺ channel inhibitors have been shown to disrupt cell proliferation (for review, see Wang, 2004). K⁺ channel activation appears to be particularly important for the early G1 phase of the cell cycle (Wonderlin & Strobl, 1996; Wang et al., 1998). The maintenance of cell membrane potential by K⁺




Fig. 2. Regulation of transport mechanisms in Jurkat T cells following stimulation of the CD95 receptor. CD95 stimulation is followed by activation of anion channels (ORCC) and, with delay, of cellular osmolyte release channels, as well as inhibition of Na^+/H^+ exchange, the K^+ channel Kv1.3 and the Ca^{2+} channels I_{CRAC} .

channels is a prerequisite for Ca^{2+} entry through I_{CRAC} (Parekh & Penner, 1997). As discussed above, I_{CRAC} activation has been shown to be required for stimulation of cell proliferation.

Reports on the role of K^+ channels in apoptosis are conflicting. In some cells, inhibition of K^+ channels appears to favour (Szabo et al., 1996, 1997, 2004; Chin et al., 1997; Miki et al., 1997; Bankers-Fulbright et al., 1998; Han et al., 2004; Pal et al., 2004; Patel & Lazdunski, 2004) and activation of K^+ channels to inhibit (Jakob & Kriegelstein, 1997; Lauritzen et al., 1997) apoptosis. Mice carrying a mutation of a G protein-coupled inward rectifier K^+ channel (*Weaver mice*) suffer from extensive neuronal cell death (Migheli et al., 1995, 1997; Murtoimaki et al., 1995; Oo et al., 1996; Harrison & Roffler-Tarlov, 1998). However, in other models, apoptosis has been reported to be stimulated by activation of K^+ channels (Yu et al., 1997; Wei et al., 2004) and to be inhibited by increase of extracellular K^+ concentration (Prehn et al., 1997; Colom et al., 1998; Lang et al., 2003e) or inhibition of K^+ channels (Gantner et al., 1995; Lang et al., 2003e). In any case, cellular




Fig. 3. Channel regulation in eryptosis. Several triggers of eryptosis (e.g., osmotic shock) stimulate the release of PGE_2 which activates Ca^{2+} permeable cation channels. The subsequent increase of cytosolic Ca^{2+} activates Ca^{2+} sensitive K^+ channels, which lead to cell shrinkage due to loss of K^+ , hyperpolarization and Cl^- exit through Cl^- channels. Ca^{2+} in addition activates a scramblase, which breaks down the phosphatidylserine asymmetry of the cell membrane. Cell shrinkage activates a phospholipase leading to formation of platelet-activating factor, sphingomyelinase activation, ceramide formation and further scramblase activation.

loss of K^+ appears to be an important trigger of apoptosis in a wide variety of cells (Beauvais et al., 1995; Benson et al., 1996; Bortner et al., 1997; Hughes, Jr. et al., 1997; Bortner & Cidlowski, 1999, 2004; Hughes, Jr. & Cidlowski, 1999; Montague et al., 1999; Gomez-Angelats et al., 2000; Maeno et al., 2000; Perez et al., 2000; Yurinskaya et al., 2005a, b). Activation of K^+ channels leads to hyperpolarization of the cell membrane, thus increasing the electrical driving force for Cl^- exit into the extracellular space. Thus, if K^+ channel activity is paralleled by Cl^- channel activity, it leads to cellular loss of KCl with osmotically obliged water and hence to cell shrinkage, a hallmark of apoptosis (Lang et al., 1998a).

As illustrated in Fig. 2, ligation of the CD95 receptor in Jurkat lymphocytes has been observed to inhibit Kv1.3 K^+ channels within a few minutes (Szabo et al., 1996, 1997, 2004). Kv1.3 is considered to be the cell volume regulatory K^+ channel of Jurkat lymphocytes (Deutsch & Chen, 1993). The channel protein is tyrosine phosphorylated upon CD95-receptor stimulation (Szabo et al., 1996; Gulbins et al., 1997) and its inhibition requires Lck⁵⁶ (Szabo et al., 1996; Gulbins et al., 1997). Similar to CD95 receptor triggering, the sphingomyelinase product ceramide inhibits Kv1.3 and induces apoptosis (Gulbins et al., 1997). Tyrosine phosphorylation of Kv1.3 has been reported to similarly inhibit channel activity in other systems (Holmes et al., 1996). Kv1.3 is stimulated by the serum- and

glucocorticoid-inducible kinase (Lang et al., 2003a), which in turn inhibits apoptosis (Aoyama et al., 2005). The early inhibition of Kv1.3 is followed by late activation of Kv1.3 upon CD95 ligation (Storey et al., 2003). The early inhibition of Kv1.3 channels following CD95 triggering prevents premature cell shrinkage, which would otherwise interfere with the signaling of apoptosis (Lang et al., 1998a). The late activation of Kv1.3 channels during the execution phase of apoptosis supports apoptotic cell shrinkage (Storey et al., 2003).

As illustrated in Fig. 3, Ca^{2+} -sensitive K^+ channels (Gardos channels) are activated by increased cytosolic Ca^{2+} activity in suicidal erythrocytes (Gardos, 1958; Grygorczyk & Schwarz, 1983; Leinders et al., 1992; Brugnara et al., 1993; Dunn, 1998; Pellegrino & Pellegrini, 1998; Shindo et al., 2000; Del Carlo et al., 2002). The activation of the channels leads to hyperpolarization and due to the high erythrocyte Cl^- permeability, to parallel exit of K^+ and Cl^- . The cellular loss of KCl and subsequent cell shrinkage stimulate eryptosis (Lang et al., 2003d). Accordingly, increase of extracellular K^+ or pharmacological inhibition of the Gardos channels blunts the cell shrinkage and apoptosis following exposure to the Ca^{2+} ionophore ionomycin (Lang et al., 2003d). The cell shrinkage following Gardos channel activation triggers the formation of platelet activating factor PAF and subsequent activation of sphingomyelinase with formation of ceramide (Lang et al., 2005c). Ceramide then potentiates the pro-apoptotic effect of Ca^{2+} (Lang et al., 2004b, 2005c).

The Role of Anion Channels, Osmolyte Transport and pH Regulation in Cell Proliferation and Apoptosis

Anion channels have been shown to be activated during cell proliferation (Shen et al., 2000; Nilius & Droogmans, 2001; Varela et al., 2004) and anion channel blockers have been shown to interfere with cell proliferation (Phipps et al., 1996; Pappas & Ritchie, 1998; Rouzair-Dubois et al., 2000; Shen et al., 2000; Wondergem et al., 2001; Jiang et al., 2004). Moreover, impaired cell proliferation has been observed in ClC_3 -deficient cells (Wang et al., 2002). Possibly, the signaling of cell proliferation needs at some stage transient cell shrinkage, which may require the activation of Cl^- channels. Usually, intracellular Cl^- is above electrochemical equilibrium and activation of Cl^- channels leads to Cl^- exit and thus depolarization. As long as K^+ channels are active, the Cl^- exit is paralleled by exit of K^+ . The loss of KCl and osmotically obliged water shrinks the cells (Lang et al., 1998a). Cell shrinkage is, for instance, required for the triggering of cytosolic Ca^{2+} oscillations in ras oncogene-expressing cells (Ritter et al.,

1993). The Ca^{2+} oscillations are in turn needed for the stimulation of cell proliferation (Fig. 1). At a later stage, proliferating cells swell due to a shift of the cell volume regulatory set point towards greater volumes and subsequent stimulation of Na^+/H^+ exchange and/or $\text{Na}^+, \text{K}^+, 2\text{Cl}^-$ cotransport (Fig. 1). At this later stage activation of Cl^- channels either tends to impede cell proliferation or maintains the activity of Na^+/H^+ exchange and/or $\text{Na}^+, \text{K}^+, 2\text{Cl}^-$ cotransport by keeping the actual cell volume below the new set point volume.

CD95 induced apoptosis of Jurkat cells (Szabo et al., 1998) and the $\text{TNF}\alpha$ - or staurosporine-induced apoptosis of various cell types (Maeno et al., 2000; Okada et al., 2004) is paralleled by activation of Cl^- channels (Fig. 2). In Jurkat cells the same channels are activated by osmotic cell swelling and are required for regulatory cell volume decrease (Lepple-Wienhues et al., 1998). In those cells activation of the Cl^- channels by cell swelling (Lepple-Wienhues et al., 1998) as well as by stimulation of the CD95 receptor (Szabo et al., 1998) requires the Src-like kinase Lck⁵⁶. The kinase is activated by ceramide (Gulbins et al., 1997), which is released by sphingomyelinase-mediated hydrolysis of sphingomyelin after stimulation of the CD95 receptor. In lymphocytes from patients with cystic fibrosis, outwardly rectifying Cl^- channels (ORCC) are resistant to activation by protein kinase A but could still be activated by cell swelling and Lck⁵⁶ (Lepple-Wienhues et al., 2001).

Activation of Cl^- channels is required for stimulation of apoptosis, which is blunted or even disrupted by Cl^- channel inhibitors. Specifically, the respective Cl^- channel blockers inhibited apoptosis in CD95-induced Jurkat cell apoptosis (Szabo et al., 1998), $\text{TNF}\alpha$ - or staurosporine-induced apoptosis of various cell types (Maeno et al., 2000; Okada et al., 2004), apoptotic death of cortical neurons (Wei et al., 2004), anti-mycin A-induced death of proximal renal tubules (Miller & Schnellmann, 1993). In addition, chloride channel blockers inhibited GABA-induced enhancement of excitotoxic cell death of rat cerebral neurons (Erdo et al., 1991); cardiomyocyte apoptosis (Takahashi et al., 2005) and eryptosis (Myssina et al., 2004).

Activation of Cl^- channels leads to cell shrinkage by triggering cellular loss of KCl (*see above*). Some anion channels further allow the permeation of organic osmolytes such as taurine (Lang et al., 2003e), which are released by cells undergoing apoptosis (Lang et al., 1998b; Moran et al., 2000). The loss of the organic osmolytes then contributes to cell shrinkage (Lang et al., 1998a). Moreover, organic osmolytes stabilize cellular proteins (for review, *see* (Lang et al., 1998a) and their loss could, indeed, destabilize proteins. For instance, inhibition of inositol uptake has been shown to induce renal failure presumably due to apoptotic death of renal tubular cells (Kitamura et al., 1998).

Many Cl^- channels further allow the passage of HCO_3^- and their activation thus leads to cytosolic acidification, a typical feature of cells entering into apoptosis (Lang et al., 2002a; Wenzel & Daniel, 2004). Acidification may promote DNA fragmentation, as DNase type II has its pH optimum in the acidic range (for review, see Shrode et al., 1997). Along those lines, CD95-induced apoptosis is accelerated by inhibition of Na^+/H^+ exchange (Lang et al., 2000a).

The Dual Role of Ion Channels – a Paradox?

At first glance it is surprising that the same or similar channels could stimulate or support both cell proliferation and apoptosis. It should be kept in mind, though, that the effect of channel activation depends on further properties of the cell. For instance, activation of K^+ channels without parallel activity of electrogenic anion transporters or Cl^- channels may hyperpolarize but not shrink the cell (Lang et al., 1998a). The influence of K^+ channel activity and cell membrane potential on cytosolic Ca^{2+} activity depends on the activity of Ca^{2+} channels.

Moreover, the temporal pattern of channel activation may be important. For instance, the functional impact of oscillating K^+ channel activity typical for proliferating cells (Pandiella et al., 1989; Lang et al., 1991) has effects different from sustained K^+ channel activation typical for apoptotic cells (Lang et al., 2003d). Oscillatory Ca^{2+} -channel activity leading to fluctuations of cytosolic Ca^{2+} concentration could depolymerize the cytoskeleton (Lang et al., 1992, 2000c; Dartsch et al., 1995; Ritter et al., 1997) but may be too short-lived to activate caspases (Whitfield et al., 1995) or the scramblase (Woon et al., 1999; Dekkers et al., 2002).

Finally, the amplitude of channel activity may be decisive for the outcome. For instance, the amplitude of TASK-3 K^+ channel activity observed during apoptosis is one order of magnitude higher than the activity of the same channels in tumor cells (Patel & Lazdunski, 2004; Wang, 2004). Similarly, the Ca^{2+} entry required for stimulation of mitogenic transcription factors may remain below the Ca^{2+} entry required for triggering of apoptosis (Whitfield et al., 1995).

Conclusions

Ion channels play an active role in the concerted cellular mechanisms leading to cell proliferation and apoptosis. They participate in the appropriate adjustment of cell volume and influence cytosolic pH and Ca^{2+} concentrations.

Typically, stimulation of cell proliferation is followed by early cell shrinkage requiring activation of Cl^- and K^+ channel activity, by cytosolic alkalization (supported by activation of the Na^+/H^+ exchanger and impeded by activation of HCO_3^- -permeable Cl^- channels), and by Ca^{2+} oscillations (requiring activation of Ca^{2+} channels). The Ca^{2+} oscillations are maintained by Ca^{2+} entry through Ca^{2+} release-activated channels I_{CRAC} , which in turn require K^+ channel-dependent maintenance of cell membrane polarization. The Ca^{2+} oscillations lead to depolymerization of the actin filament network, with subsequent disinhibition of Na^+/H^+ exchanger and/or $\text{Na}^+,\text{K}^+,\text{2Cl}^-$ resulting in cell swelling.

Typically, apoptosis eventually leads to cell shrinkage due to activation of K^+ and/or Cl^- channels and organic osmolyte release. The activation of Cl^- channels and inhibition of Na^+/H^+ exchangers leads to cytosolic acidification. While I_{CRAC} is inhibited during CD95-induced apoptosis, sustained Ca^{2+} entry through Ca^{2+} -permeable cation channels is able to trigger apoptosis.

Several channels play dual roles in both cell proliferation and apoptosis. The effect of channel activation critically depends on the temporal pattern and amplitude of channel activity as well as the interplay with other channels, transporters and signaling pathways.

Despite the generation of a tremendous body of experimental evidence we are still far from fully understanding the complex interplay between channel activity and signaling of proliferating or dying cells. Clearly, further experimental efforts are needed to address the many open questions.

The authors acknowledge the meticulous preparation of the manuscript by Lejla Subasic. The work of the authors was supported by the Deutsche Forschungsgemeinschaft, Nr. La 315/4-3, La 315/6-1, Le 792/3-3, DFG Schwerpunkt Intrazelluläre Lebensformen La 315/11-1, and Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (Center for Interdisciplinary Clinical Research) 01 KS 9602.

References

- Adams, T.E., McKern, N.M., Ward, C.W. 2004. Signalling by the type 1 insulin-like growth factor receptor: interplay with the epidermal growth factor receptor. *Growth Factors* **22**:89–95
- Albright, C.D., da Costa, K.A., Craciunescu, C.N., Klem, E., Mar, M.H., Zeisel, S.H. 2005. Regulation of choline deficiency apoptosis by epidermal growth factor in CWSV-1 rat hepatocytes. *Cell Physiol. Biochem.* **15**:59–68
- Alisi, A., Demori, I., Spagnuolo, S., Pierantozzi, E., Fugassa, E., Leoni, S. 2005. Thyroid status affects rat liver regeneration after partial hepatectomy by regulating cell cycle and apoptosis. *Cell Physiol. Biochem.* **15**:69–76
- Aoyama, T., Matsui, T., Novikov, M., Park, J., Hemmings, B., Rosenzweig, A. 2005. Serum and glucocorticoid-responsive kinase-1 regulates cardiomyocyte survival and hypertrophic response. *Circulation* **111**:1652–1659

- Bankers-Fulbright, J.L., Kephart, G.M., Loegering, D.A., Bradford, A.L., Okada, S., Kita, H., Gleich, G.J. 1998. Sulfonylureas inhibit cytokine-induced eosinophil survival and activation. *J. Immunol.* **160**:5546–5553
- Barvitenko, N.N., Adragna, N.C., Weber, R.E. 2005. Erythrocyte signal transduction pathways, their oxygenation dependence and functional significance. *Cell Physiol. Biochem.* **15**:1–18
- Beauvais, F., Michel, L., Dubertret, R. 1995. Human eosinophils in culture undergo a striking and rapid shrinkage during apoptosis. Role of K⁺ channels. *J. Leukoc. Biol.* **57**:851–855
- Bennekou, P. 1993. The voltage-gated non-selective cation channel from human red cells is sensitive to acetylcholine. *Biochim. Biophys. Acta* **1147**:165–167
- Benson, R.S., Heer, S., Dive, C., Watson, A.J. 1996. Characterization of cell volume loss in CEM-C7A cells during dexamethasone-induced apoptosis. *Am. J. Physiol.* **270**:C1190–C1203
- Berg, C.P., Engels, I.H., Rothbart, A., Lauber, K., Renz, A., Schlosser, S.F., Schulze-Osthoff, K., Wesselborg, S. 2001. Human mature red blood cells express caspase-3 and caspase-8, but are devoid of mitochondrial regulators of apoptosis. *Cell Death. Differ.* **8**:1197–1206
- Bernhardt, I., Hall, A.C., Ellory, J.C. 1991. Effects of low ionic strength media on passive human red cell monovalent cation transport. *J. Physiol.* **434**:489–506
- Berridge, M.J., Bootman, M.D., Lipp, P. 1998. Calcium—a life and death signal. *Nature* **395**:645–648
- Berridge, M.J., Bootman, M.D., Roderick, H.L. 2003. Calcium signalling: dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell. Biol.* **4**:517–529
- Berridge, M.J., Lipp, P., Bootman, M.D. 2000. The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* **1**:11–21
- Bikfalvi, A., Savona, C., Perollet, C., Javerzat, S. 1998. New insights in the biology of fibroblast growth factor-2. *Angiogenesis.* **1**:155–173
- Bilmen, S., Aksu, T.A., Gumuslu, S., Korgun, D.K., Canatan, D. 2001. Antioxidant capacity of G-6-PD-deficient erythrocytes. *Clin. Chim. Acta* **303**:83–86
- Boas, F.E., Forman, L., Beutler, E. 1998. Phosphatidylserine exposure and red cell viability in red cell aging and in hemolytic anemia. *Proc. Natl. Acad. Sci. USA* **95**:3077–3081
- Bortner, C.D., Cidlowski, J.A. 1998. A necessary role for cell shrinkage in apoptosis. *Biochem. Pharmacol.* **56**:1549–1559
- Bortner, C.D., Cidlowski, J.A. 1999. Caspase independent/dependent regulation of K(+), cell shrinkage, and mitochondrial membrane potential during lymphocyte apoptosis. *J. Biol. Chem.* **274**:21953–21962
- Bortner, C.D., Cidlowski, J.A. 2004. The role of apoptotic volume decrease and ionic homeostasis in the activation and repression of apoptosis. *Pfluegers Arch.* **448**:313–318
- Bortner, C.D., Hughes, F.M., Jr., Cidlowski, J.A. 1997. A primary role for K⁺ and Na⁺ efflux in the activation of apoptosis. *J. Biol. Chem.* **272**:32436–32442
- Bosman, G.J.C.G.M., Willekens, F.L.A. 2005. Erythrocyte aging: A more than superficial resemblance to apoptosis? *Cell Physiol. Biochem.* **16**:1–8
- Brand, V.B., Sandu, C.D., Duranton, C., Tanneur, V., Lang, K.S., Huber, S.M., Lang, F. 2003. Dependence of Plasmodium falciparum *in vitro* growth on the cation permeability of the human host erythrocyte. *Cell Physiol. Biochem.* **13**:347–356
- Bratosin, D., Leszczynski, S., Sartiaux, C., Fontaine, O., Descamps, J., Huart, J.J., Poplineau, J., Goudaliez, F., Aminoff, D., Montreuil, J. 2001. Improved storage of erythrocytes by prior leukodepletion: flow cytometric evaluation of stored erythrocytes. *Cytometry* **46**:351–356
- Brugnara, C., Franceschi, L., Alper, S.L. 1993. Inhibition of Ca(2+)-dependent K⁺ transport and cell dehydration in sickle erythrocytes by clotrimazole and other imidazole derivatives. *J. Clin. Invest.* **92**:520–526
- Cabado, A.G., Vieytes, M.R., Botana, L.M. 1994. Effect of ion composition on the changes in membrane potential induced with several stimuli in rat mast cells. *J. Cell Physiol.* **158**:309–316
- Cariers, A., Reinehr, R., Fischer, R., Warskulat, U., Haussinger, D. 2002. c-Jun-N-terminal kinase dependent membrane targeting of CD95 in rat hepatic stellate cells. *Cell Physiol. Biochem.* **12**:179–186
- Chan, H.C., Goldstein, J., Nelson, D.J. 1992. Alternate pathways for chloride conductance activation in normal and cystic fibrosis airway epithelial cells. *Am. J. Physiol.* **262**:C1273–C1283
- Chin, L.S., Park, C.C., Zitnay, K.M., Sinha, M., DiPatri, A.J., Jr., Perillan, P., Simard, J.M. 1997. 4-Aminopyridine causes apoptosis and blocks an outward rectifier K⁺ channel in malignant astrocytoma cell lines. *J. Neurosci. Res.* **48**:122–127
- Christophersen, P., Bennekou, P. 1991. Evidence for a voltage-gated, non-selective cation channel in the human red cell membrane. *Biochim. Biophys. Acta* **1065**:103–106
- Colom, L.V., Diaz, M.E., Beers, D.R., Neely, A., Xie, W.J., Appel, S.H. 1998. Role of potassium channels in amyloid-induced cell death. *J. Neurochem.* **70**:1925–1934
- Dangel, G.R., Lang, F., Lepple-Wienhues, A. 2005. Effect of Sphingosin on Ca²⁺ entry and mitochondrial potential of Jurkat T cells – interaction with Bcl2. *Cell Physiol. Biochem.* **16**:9–14
- Dartsch, P.C., Ritter, M., Gschwentner, M., Lang, H.J., Lang, F. 1995. Effects of calcium channel blockers on NIH 3T3 fibroblasts expressing the Ha-ras oncogene. *Eur. J. Cell. Biol.* **67**:372–378
- Daugas, E., Cande, C., Kroemer, G. 2001. Erythrocytes: death of a mummy. *Cell Death. Differ.* **8**:1131–1133
- Davies, A.M. 2003. Regulation of neuronal survival and death by extracellular signals during development. *EMBO J.* **22**:2537–2545
- DeCoursey, T.E., Chandy, K.G., Gupta, S., Cahalan, M.D. 1984. Voltage-gated K⁺ channels in human T lymphocytes: a role in mitogenesis? *Nature* **307**:465–468
- Dekkers, D.W., Comfurius, P., Bevers, E.M., Zwaal, R.F. 2002. Comparison between Ca²⁺-induced scrambling of various fluorescently labelled lipid analogues in red blood cells. *Biochem. J.* **362**:741–747
- Del Carlo, B., Del, P., Pellegrini, M., Pellegrino, M. 2002. Calmodulin antagonists do not inhibit IK(Ca) channels of human erythrocytes. *Biochim. Biophys. Acta* **1558**:133–141
- Deutsch, C., Chen, L.Q. 1993. Heterologous expression of specific K⁺ channels in T lymphocytes: functional consequences for volume regulation. *Proc. Natl. Acad. Sci. USA* **90**:10036–10040
- Dinudom, A., Komwatana, P., Young, J.A., Cook, D.I. 1995. Control of the amiloride-sensitive Na⁺ current in mouse salivary ducts by intracellular anions is mediated by a G protein. *J. Physiol.* **487**:549–555
- Dunn, P.M. 1998. The action of blocking agents applied to the inner face of Ca(2+)-activated K⁺ channels from human erythrocytes. *J. Membrane Biol.* **165**:133–143
- Duranton, C., Huber, S., Tanneur, V., Lang, K., Brand, V., Sandu, C., Lang, F. 2003. Electrophysiological properties of the Plasmodium falciparum-induced cation conductance of human erythrocytes. *Cell Physiol. Biochem.* **13**:189–198
- Duranton, C., Huber, S.M., Lang, F. 2002. Oxidation induces a Cl(–)-dependent cation conductance in human red blood cells. *J. Physiol.* **539**:847–855

- Enomoto, K., Cossu, M.F., Edwards, C., Oka, T. 1986. Induction of distinct types of spontaneous electrical activities in mammary epithelial cells by epidermal growth factor and insulin. *Proc Natl. Acad. Sci. USA* **83**:4754–4758
- Erdo, S., Michler, A., Wolff, J.R. 1991. GABA accelerates excitotoxic cell death in cortical cultures: protection by blockers of GABA-gated chloride channels. *Brain Res.* **542**:254–258
- Fadok, V.A., Bratton, D.L., Rose, D.M., Pearson, A., Ezekewitz, R.A., Henson, P.M. 2000. A receptor for phosphatidylserine-specific clearance of apoptotic cells. *Nature* **405**:85–90
- Faehling, M., Koch, E.D., Raithe, J., Trischler, G., Waltenberger, J. 2001. Vascular endothelial growth factor-A activates Ca^{2+} -activated K^{+} channels in human endothelial cells in culture. *Int. J. Biochem. Cell Biol.* **33**:337–346
- Fillon, S., Klingel, K., Warntges, S., Sauter, M., Gabrysch, S., Pestel, S., Tanneur, V., Waldegger, S., Zipfel, A., Viebahn, R., Haussinger, D., Broer, S., Kandolf, R., Lang, F. 2002. Expression of the serine/threonine kinase hSGK1 in chronic viral hepatitis. *Cell Physiol. Biochem.* **12**:47–54
- Gamper, N., Huber, S.M., Badawi, K., Lang, F. 2000. Cell volume-sensitive sodium channels upregulated by glucocorticoids in U937 macrophages. *Pfluegers Arch.* **441**:281–286
- Gantner, F., Uhlig, S., Wendel, A. 1995. Quinine inhibits release of tumor necrosis factor, apoptosis, necrosis and mortality in a murine model of septic liver failure. *Eur. J. Pharmacol.* **294**:353–355
- Gardos, G. 1958. The function of calcium in the potassium permeability of human erythrocytes. *Biochim. Biophys. Acta* **30**:653–654
- Gomez-Angelats, M., Bortner, C.D., Cidlowski, J.A. 2000. Protein kinase C (PKC) inhibits fas receptor-induced apoptosis through modulation of the loss of K^{+} and cell shrinkage. A role for PKC upstream of caspases. *J. Biol. Chem.* **275**:19609–19619
- Green, D.R., Reed, J.C. 1998. Mitochondria and apoptosis. *Science* **281**:1309–1312
- Grygorczyk, R., Schwarz, W. 1983. Properties of the Ca^{2+} -activated K^{+} conductance of human red cells as revealed by the patch-clamp technique. *Cell Calcium* **4**:499–510
- Gulbins, E., Jekle, A., Ferlinz, K., Grassme, H., Lang, F. 2000. Physiology of apoptosis. *Am. J. Physiol.* **279**:F605–F615
- Gulbins, E., Szabo, I., Baltzer, K., Lang, F. 1997. Ceramide-induced inhibition of T lymphocyte voltage-gated potassium channel is mediated by tyrosine kinases. *Proc. Natl. Acad. Sci. USA* **94**:7661–7666
- Han, H., Wang, J., Zhang, Y., Long, H., Wang, H., Xu, D., Wang, Z. 2004. HERG K channel conductance promotes H_2O_2 -induced apoptosis in HEK293 cells: cellular mechanisms. *Cell. Physiol. Biochem.* **14**:121–134
- Harrison, S.M., Roffler-Tarlov, S.K. 1998. Cell death during development of testis and cerebellum in the mutant mouse weaver. *Dev. Biol.* **195**:174–186
- Henson, P.M., Bratton, D.L., Fadok, V.A. 2001. The phosphatidylserine receptor: a crucial molecular switch? *Nat. Rev. Mol. Cell Biol.* **2**:627–633
- Holmes, T.C., Fadool, D.A., Levitan, I.B. 1996. Tyrosine phosphorylation of the Kv1.3 potassium channel. *J. Neurosci.* **16**:1581–1590
- Huber, S.M., Gamper, N., Lang, F. 2001. Chloride conductance and volume-regulatory nonselective cation conductance in human red blood cell ghosts. *Pfluegers Arch.* **441**:551–558
- Hughes, F.M., Jr., Bortner, C.D., Purdy, G.D., Cidlowski, J.A. 1997. Intracellular K^{+} suppresses the activation of apoptosis in lymphocytes. *J. Biol. Chem.* **272**:30567–30576
- Hughes, F.M., Jr., Cidlowski, J.A. 1999. Potassium is a critical regulator of apoptotic enzymes *in vitro* and *in vivo*. *Adv. Enzyme Regul.* **39**:157–171
- Jakob, R., Krieglstein, J. 1997. Influence of flupirtine on a G-protein coupled inwardly rectifying potassium current in hippocampal neurones. *Br. J. Pharmacol.* **122**:1333–1338
- Jiang, B., Hattori, N., Liu, B., Nakayama, Y., Kitagawa, K., Inagaki, C. 2004. Suppression of cell proliferation with induction of p21 by Cl^{-} channel blockers in human leukemic cells. *Eur. J. Pharmacol.* **488**:27–34
- Jones, G.S., Knauf, P.A. 1985. Mechanism of the increase in cation permeability of human erythrocytes in low-chloride media. Involvement of the anion transport protein capnophorin. *J. Gen. Physiol.* **86**:721–738
- Kaestner, L., Bollensdorff, C., Bernhardt, I. 1999. Non-selective voltage-activated cation channel in the human red blood cell membrane. *Biochim. Biophys. Acta* **1417**:9–15
- Kitamura, H., Yamauchi, A., Sugiura, T., Matsuoka, Y., Horio, M., Tohyama, M., Shimada, S., Imai, E., Hori, M. 1998. Inhibition of myo-inositol transport causes acute renal failure with selective medullary injury in the rat. *Kidney Int.* **53**:146–153
- Koch, J., Korbmacher, C. 1999. Osmotic shrinkage activates nonselective cation (NSC) channels in various cell types. *J. Membrane Biol.* **168**:131–139
- Kohn, K.W., Pommier, Y. 2005. Molecular interaction map of the p53 and Mdm2 logic elements, which control the Off-On switch of p53 in response to DNA damage. *Biochem. Biophys. Res. Commun.* **331**:816–827
- LaCelle, P.L., Rothstein, A. 1966. The passive permeability of the red blood cell in cations. *J. Gen. Physiol.* **50**:171–188
- Lang, F., Busch, G.L., Ritter, M., Volk, H., Waldegger, S., Gulbins, E., Haussinger, D. 1998a. Functional significance of cell volume regulatory mechanisms. *Physiol. Rev.* **78**:247–306
- Lang, K.S., Duranton, C., Poehlmann, H., Myssina, S., Bauer, C., Lang, F., Wieder, T., Huber, S.M. 2003b. Cation channels trigger apoptotic death of erythrocytes. *Cell Death Differ.* **10**:249–256
- Lang, K.S., Fillon, S., Schneider, D., Rammensee, H.G., Lang, F. 2002a. Stimulation of TNF alpha expression by hyperosmotic stress. *Pfluegers Arch.* **443**:798–803
- Lang, F., Friedrich, F., Kahn, E., Woll, E., Hammerer, M., Waldegger, S., Maly, K., Grunicke, H. 1991. Bradykinin-induced oscillations of cell membrane potential in cells expressing the Ha-ras oncogene. *J. Biol. Chem.* **266**:4938–4942
- Lang, F., Henke, G., Embark, H.M., Waldegger, S., Palmada, M., Bohmer, C., Vallon, V. 2003a. Regulation of channels by the serum and glucocorticoid-inducible kinase-implications for transport, excitability and cell proliferation. *Cell Physiol. Biochem.* **13**:41–50
- Lang, P.A., Kaiser, S., Myssina, S., Wieder, T., Lang, F., Huber, S.M. 2003d. Role of Ca^{2+} -activated K^{+} channels in human erythrocyte apoptosis. *Am. J. Physiol.* **285**:C1553–C1560
- Lang, P.A., Kempe, D.S., Myssina, S., Tanneur, V., Birka, C., Laufer, S., Lang, F., Wieder, T., Huber, S.M. 2005b. PGE(2) in the regulation of programmed erythrocyte death. *Cell Death Differ.* **12**:415–428
- Lang, P.A., Kempe, D.S., Tanneur, V., Eisele, K., Klarl, B.A., Myssina, S., Jendrosseck, V., Ishii, S., Shimizu, T., Waidmann, M., Hessler, G., Huber, S.M., Lang, F., Wieder, T. 2005c. Stimulation of erythrocyte ceramide formation by platelet-activating factor. *J. Cell Sci.* **118**:1233–1243
- Lang, K.S., Lang, P.A., Bauer, C., Duranton, C., Wieder, T., Huber, S.M., Lang, F. 2005a. Mechanisms of suicidal erythrocyte death. *Cell Physiol. Biochem.* **15**:195–202
- Lang, F., Lang, P.A., Lang, K.S., Brand, V., Tanneur, V., Duranton, C., Wieder, T., Huber, S.M. 2004a. Channel-induced apoptosis of infected host cells-the case of malaria. *Pfluegers Arch.* **448**:319–324

- Lang, F., Madlung, J., Bock, J., Lukewille, U., Kaltenbach, S., Lang, K.S., Belka, C., Wagner, C.A., Lang, H.J., Gulbins, E., Lepple-Wienhues, A. 2000a. Inhibition of Jurkat-T-lymphocyte Na^+/H^+ -exchanger by CD95(Fas/Apo-1)-receptor stimulation. *Pfluegers Arch.* **440**:902–907
- Lang, F., Madlung, J., Siemen, D., Ellory, C., Lepple-Wienhues, A., Gulbins, E. 2000b. The involvement of caspases in the CD95(Fas/Apo-1)- but not swelling-induced cellular taurine release from Jurkat T-lymphocytes. *Pfluegers Arch.* **440**:93–99
- Lang, F., Madlung, J., Uhlemann, A.C., Risler, T., Gulbins, E. 1998b. Cellular taurine release triggered by stimulation of the Fas(CD95) receptor in Jurkat lymphocytes. *Pfluegers Arch.* **436**:377–383
- Lang, K.S., Myssina, S., Brand, V., Sandu, C., Lang, P.A., Berchtold, S., Huber, S.M., Lang, F., Wieder, T. 2004b. Involvement of ceramide in hyperosmotic shock-induced death of erythrocytes. *Cell Death Differ.* **11**:231–243
- Lang, K.S., Myssina, S., Tanneur, V., Wieder, T., Huber, S.M., Lang, F., Duranton, C. 2003c. Inhibition of erythrocyte cation channels and apoptosis by ethylisopropylamiloride. *Naunyn Schmiedeberg's Arch. Pharmacol.* **367**:391–396
- Lang, F., Ritter, M., Gamper, N., Huber, S., Fillon, S., Tanneur, V., Lepple-Wienhues, A., Szabo, I., Gulbins, E. 2000c. Cell volume in the regulation of cell proliferation and apoptotic cell death. *Cell Physiol. Biochem.* **10**:417–428
- Lang, K.S., Roll, B., Myssina, S., Schittenhelm, M., Scheel-Walter, H.G., Kanz, L., Fritz, J., Lang, F., Huber, S.M., Wieder, T. 2002b. Enhanced erythrocyte apoptosis in sickle cell anemia, thalassemia and glucose-6-phosphate dehydrogenase deficiency. *Cell Physiol. Biochem.* **12**:365–372
- Lang, F., Szabo, I., Lepple-Wienhues, A., Siemen, D., Gulbins, E. 1999. Physiology of Receptor-Mediated Lymphocyte Apoptosis. *News Physiol. Sci.* **14**:194–200
- Lang, F., Waldegger, S., Woell, E., Ritter, M., Maly, K., Grunicke, H. 1992. Effects of inhibitors and ion substitutions on oscillations of cell membrane potential in cells expressing the RAS oncogene. *Pfluegers Arch.* **421**:416–424
- Lang, P.A., Warskulat, U., Heller-Stilb, B., Huang, D.Y., Grenz, A., Myssina, S., Duszenko, M., Lang, F., Haussinger, D., Vallon, V., Wieder, T. 2003e. Blunted apoptosis of erythrocytes from taurine transporter deficient mice. *Cell Physiol. Biochem.* **13**:337–346
- Lauritzen, I., Weille, J.R., Lazdunski, M. 1997. The potassium channel opener (–)-cromakalim prevents glutamate-induced cell death in hippocampal neurons. *J. Neurochem.* **69**:1570–1579
- Leinders, T., Kleef, R.G., Vijverberg, H.P. 1992. Single Ca^{2+} -activated K^+ channels in human erythrocytes: Ca^{2+} dependence of opening frequency but not of open lifetimes. *Biochim. Biophys. Acta* **1112**:67–74
- Lepple-Wienhues, A., Belka, C., Laun, T., Jekle, A., Walter, B., Wieland, U., Welz, M., Heil, L., Kun, J., Busch, G., Weller, M., Bamberg, M., Gulbins, E., Lang, F. 1999. Stimulation of CD95 (Fas) blocks T lymphocyte calcium channels through sphingomyelinase and sphingolipids. *Proc Natl. Acad. Sci. USA* **96**:13795–13800
- Lepple-Wienhues, A., Szabo, I., Laun, T., Kaba, N.K., Gulbins, E., Lang, F. 1998. The tyrosine kinase p56lck mediates activation of swelling-induced chloride channels in lymphocytes. *J. Cell Biol.* **141**:281–286
- Lepple-Wienhues, A., Wieland, U., Laun, T., Heil, L., Stern, M., Lang, F. 2001. A src-like kinase activates outwardly rectifying chloride channels in CFTR-defective lymphocytes. *FASEB J.* **15**:927–931
- Liu, X.H., Kirschenbaum, A., Yu, K., Yao, S., Levine, A.C. 2005. Cyclooxygenase-2 suppresses hypoxia-induced apoptosis via a combination of direct and indirect inhibition of p53 activity in a human prostate cancer cell line. *J. Biol. Chem.* **280**:3817–3823
- Liu, X.M., Tao, M., Han, X.D., Fan, Q., Lin, J.R. 2001. Gating kinetics of potassium channel and effects of nerve growth factors in PC12 cells analyzed with fractal model. *Acta Pharmacol. Sin.* **22**:103–110
- Long, H., Han, H., Yang, B., Wang, Z. 2003. Opposite cell density-dependence between spontaneous and oxidative stress-induced apoptosis in mouse fibroblast L-cells. *Cell Physiol. Biochem.* **13**:401–414
- Maeno, E., Ishizaki, Y., Kanaseki, T., Hazama, A., Okada, Y. 2000. Normotonic cell shrinkage because of disordered volume regulation is an early prerequisite to apoptosis. *Proc. Natl. Acad. Sci. USA* **97**:9487–9492
- Marunaka, Y., Nakahara, T., Tohda, H. 1994. Cytosolic $[\text{Cl}^-]$ regulates Na^+ absorption in fetal alveolar epithelium?: roles of cAMP and Cl^- channels. *Jpn. J. Physiol.* **44**:S281–S288
- Mauro, T., Dixon, D.B., Komuves, L., Hanley, K., Pappone, P.A. 1997. Keratinocyte K^+ channels mediate Ca^{2+} -induced differentiation. *J. Invest. Dermatol.* **108**:864–870
- Mavelli, I., Ciriolo, M.R., Rossi, L., Meloni, T., Forteleoni, G., De Flora, A. De , Benatti, U., Morelli, A., Rotilio, G. 1984. Favism: a hemolytic disease associated with increased superoxide dismutase and decreased glutathione peroxidase activities in red blood cells. *Eur. J. Biochem.* **139**:13–18
- Michea, L., Ferguson, D.R., Peters, E.M., Andrews, P.M., Kirby, M.R., Burg, M.B. 2000. Cell cycle delay and apoptosis are induced by high salt and urea in renal medullary cells. *Am. J. Physiol.* **278**:F209–F218
- Migheli, A., Attanasio, A., Lee, W.H., Bayer, S.A., Ghatti, B. 1995. Detection of apoptosis in weaver cerebellum by electron microscopic *in situ* end-labeling of fragmented DNA. *Neurosci. Lett.* **199**:53–56
- Migheli, A., Piva, R., Wei, J., Attanasio, A., Casolino, S., Hodes, M.E., Dlouhy, S.R., Bayer, S.A., Ghatti, B. 1997. Diverse cell death pathways result from a single missense mutation in weaver mouse. *Am. J. Pathol.* **151**:1629–1638
- Miki, T., Tashiro, F., Iwanaga, T., Nagashima, K., Yoshitomi, H., Aihara, H., Nitta, Y., Gono, T., Inagaki, N., Miyazaki, J., Seino, S. 1997. Abnormalities of pancreatic islets by targeted expression of a dominant-negative KATP channel. *Proc Natl. Acad. Sci. USA* **94**:11969–11973
- Miller, G.W., Schnellmann, R.G. 1993. Cytoprotection by inhibition of chloride channels: the mechanism of action of glycine and strychnine. *Life Sci.* **53**:1211–1215
- Montague, J.W., Bortner, C.D., Hughes, F.M., Jr., Cidlowski, J.A. 1999. A necessary role for reduced intracellular potassium during the DNA degradation phase of apoptosis. *Steroids* **64**:563–569
- Moran, J., Hernandez-Pech, X., Merchant-Larios, H., Pasantes-Morales, H. 2000. Release of taurine in apoptotic cerebellar granule neurons in culture. *Pfluegers Arch.* **439**:271–277
- Murtomaki, S., Trenkner, E., Wright, J.M., Saksela, O., Liesi, P. 1995. Increased proteolytic activity of the granule neurons may contribute to neuronal death in the weaver mouse cerebellum. *Dev. Biol.* **168**:635–648
- Myssina, S., Lang, P.A., Kempe, D.S., Kaiser, S., Huber, S.M., Wieder, T., Lang, F. 2004. Cl^- channel blockers NPPB and niflumic acid blunt Ca^{2+} -induced erythrocyte 'apoptosis'. *Cell Physiol. Biochem.* **14**:241–248
- Nilius, B., Droogmans, G. 2001. Ion channels and their functional role in vascular endothelium. *Physiol. Rev.* **81**:1415–1459
- Nilius, B., Wohlrab, W. 1992. Potassium channels and regulation of proliferation of human melanoma cells. *J. Physiol.* **445**:537–548

- O'Laigue, P.H., Huttner, S.L., Vandenberg, C.A., Morrison-Graham, K., Horn, R. 1985. Morphological properties and membrane channels of the growth cones induced in PC12 cells by nerve growth factor. *J. Neurosci. Res.* **13**:301–321
- Okada, Y., Maeno, E., Shimizu, T., Manabe, K., Mori, S., Nabekura, T. 2004. Dual roles of plasmalemmal chloride channels in induction of cell death. *Pfluegers Arch.* **448**:287–295
- Oo, T.F., Blazeski, R., Harrison, S.M., Henchcliffe, C., Mason, C.A., Roffler-Tarlov, S.K., Burke, R.E. 1996. Neuron death in the substantia nigra of weaver mouse occurs late in development and is not apoptotic. *J. Neurosci.* **16**:6134–6145
- Pal, S., He, K., Aizenman, E. 2004. Nitrosative stress and potassium channel-mediated neuronal apoptosis: is zinc the link? *Pfluegers Arch.* **448**:296–303
- Pandiella, A., Magni, M., Lovisolo, D., Meldolesi, J. 1989. The effect of epidermal growth factor on membrane potential. Rapid hyperpolarization followed by persistent fluctuations. *J. Biol. Chem.* **264**:12914–12921
- Pappas, C.A., Ritchie, J.M. 1998. Effect of specific ion channel blockers on cultured Schwann cell proliferation. *Glia* **22**:113–120
- Papnone, P.A., Ortiz-Miranda, S.I. 1993. Blockers of voltage-gated K channels inhibit proliferation of cultured brown fat cells. *Am. J. Physiol.* **264**:C1014–C1019
- Parekh, A.B., Penner, R. 1997. Store depletion and calcium influx. *Physiol. Rev.* **77**:901–930
- Parekh, A.B., Putney, J.W. Jr. 2005. Store-operated calcium channels. *Physiol. Rev.* **85**:757–810
- Patel, A.J., Lazdunski, M. 2004. The 2P-domain K⁺ channels: role in apoptosis and tumorigenesis. *Pfluegers Arch.* **448**:261–273
- Pellegrino, M., Pellegrini, M. 1998. Modulation of Ca²⁺-activated K⁺ channels of human erythrocytes by endogenous cAMP-dependent protein kinase. *Pfluegers Arch.* **436**:749–756
- Perez, G.I., Maravei, D.V., Trbovich, A.M., Cidlowski, J.A., Tilly, J.L., Hughes, F.M. Jr. 2000. Identification of potassium-dependent and -independent components of the apoptotic machinery in mouse ovarian germ cells and granulosa cells. *Biol. Reprod.* **63**:1358–1369
- Phipps, D.J., Branch, D.R., Schlichter, L.C. 1996. Chloride-channel block inhibits T lymphocyte activation and signalling. *Cell Signal.* **8**:141–149
- Pozzi, S., Malferrari, G., Biunno, I., Samaja, M. 2002. Low-flow ischemia and hypoxia stimulate apoptosis in perfused hearts independently of reperfusion. *Cell Physiol. Biochem.* **12**:39–46
- Prehn, J.H., Jordan, J., Ghadge, G.D., Preis, E., Galindo, M.F., Roos, R.P., Kriegelstein, J., Miller, R.J. 1997. Ca²⁺ and reactive oxygen species in staurosporine-induced neuronal apoptosis. *J. Neurochem.* **68**:1679–1685
- Qian, D., Weiss, A. 1997. T cell antigen receptor signal transduction. *Curr. Opin. Cell Biol.* **9**:205–212
- Rice, L., Alfrey, C.P. 2005. The negative regulation of red cell mass by neocytolysis: physiologic and pathophysiologic manifestations. *Cell Physiol. Biochem.* **15**:245–250
- Ritter, M., Woll, E., Haller, T., Dartsch, P.C., Zwierzina, H., Lang, F. 1997. Activation of Na⁺/H⁺-exchanger by transforming Ha-ras requires stimulated cellular calcium influx and is associated with rearrangement of the actin cytoskeleton. *Eur. J. Cell Biol.* **72**:222–228
- Ritter, M., Woll, E., Waldegger, S., Haussinger, D., Lang, H.J., Scholz, W., Scholkens, B., Lang, F. 1993. Cell shrinkage stimulates bradykinin-induced cell membrane potential oscillations in NIH 3T3 fibroblasts expressing the ras-oncogene. *Pfluegers Arch.* **423**:221–224
- Rosette, C., Karin, M. 1996. Ultraviolet light and osmotic stress: activation of the JNK cascade through multiple growth factor and cytokine receptors. *Science* **274**:1194–1197
- Rotoli, B.M., Uggeri, J., Dall'Asta, V., Visigalli, R., Barilli, A., Gatti, R., Orlandini, G., Gazzola, G.C., Bussolati, O. 2005. Inhibition of glutamine synthetase triggers apoptosis in asparaginase-resistant cells. *Cell Physiol. Biochem.* **15**:281–292
- Rouzaire-Dubois, B., Milandri, J.B., Bostel, S., Dubois, J.M. 2000. Control of cell proliferation by cell volume alterations in rat C6 glioma cells. *Pfluegers Arch.* **440**:881–888
- Sanders, D.A., Fiddes, I., Thompson, D.M., Philpott, M.P., Westgate, G.E., Kealey, T. 1996. In the absence of streptomycin, minoxidil potentiates the mitogenic effects of fetal calf serum, insulin-like growth factor 1, and platelet-derived growth factor on NIH 3T3 fibroblasts in a K⁺ channel-dependent fashion. *J. Invest. Dermatol.* **107**:229–234
- Santella, L. 1998. The role of calcium in the cell cycle: facts and hypotheses. *Biochem Biophys. Res. Commun.* **244**:317–324
- Santella, L., Kyozuka, K., De Riso, L., De, Carafoli, E. 1998. Calcium, protease action, and the regulation of the cell cycle. *Cell Calcium* **23**:123–130
- Shen, M.R., Droogmans, G., Eggermont, J., Voets, T., Ellory, J.C., Nilius, B. 2000. Differential expression of volume-regulated anion channels during cell cycle progression of human cervical cancer cells. *J. Physiol.* **529 Pt 2**:385–394
- Shindo, M., Imai, Y., Sohma, Y. 2000. A novel type of ATP block on a Ca²⁺-activated K⁺ channel from bullfrog erythrocytes. *Biophys J* **79**:287–297
- Shrode, L.D., Tapper, H., Grinstein, S. 1997. Role of intracellular pH in proliferation, transformation, and apoptosis. *J. Bioenerg. Biomembr.* **29**:393–399
- Skryma, R.N., Prevarkaya, N.B., Dufy-Barbe, L., Odessa, M.F., Audin, J., Dufy, B. 1997. Potassium conductance in the androgen-sensitive prostate cancer cell line, LNCaP: involvement in cell proliferation. *Prostate* **33**:112–122
- Spasova, M.A., Soboloff, J., He, L.P., Hewavitharana, T., Xu, W., Venkatachalam, K., van Rossum, D.B. van, Patterson, R.L., Gill, D.L. 2004. Calcium entry mediated by SOCs and TRP channels: variations and enigma. *Biochim. Biophys. Acta* **1742**:9–20
- Storey, N.M., Gomez-Angelats, M., Bortner, C.D., Armstrong, D.L., Cidlowski, J.A. 2003. Stimulation of Kv1.3 potassium channels by death receptors during apoptosis in Jurkat T lymphocytes. *J. Biol. Chem.* **278**:33319–33326
- Strobl, J.S., Wonderlin, W.F., Flynn, D.C. 1995. Mitogenic signal transduction in human breast cancer cells. *Gen. Pharmacol.* **26**:1643–1649
- Sturm, J.W., Zhang, H., Magdeburg, R., Hasenberg, T., Bonninghoff, R., Oulmi, J., Keese, M., McCuskey, R. 2004. Altered apoptotic response and different liver structure during liver regeneration in FGF-2-deficient mice. *Cell Physiol. Biochem.* **14**:249–260
- Szabo, I., Adams, C., Gulbins, E. 2004. Ion channels and membrane rafts in apoptosis. *Pfluegers Arch.* **448**:304–312
- Szabo, I., Gulbins, E., Apfel, H., Zhang, X., Barth, P., Busch, A.E., Schlottmann, K., Pongs, O., Lang, F. 1996. Tyrosine phosphorylation-dependent suppression of a voltage-gated K⁺ channel in T lymphocytes upon Fas stimulation. *J. Biol. Chem.* **271**:20465–20469
- Szabo, I., Gulbins, E., Lang, F. 1997. Regulation of Kv1.3 during Fas-induced apoptosis. *Cell Physiol. Biochem.* **7**:148–158
- Szabo, I., Lepple-Wienhues, A., Kaba, K.N., Zoratti, M., Gulbins, E., Lang, F. 1998. Tyrosine kinase-dependent activation of a chloride channel in CD95-induced apoptosis in T lymphocytes. *Proc. Natl. Acad. Sci. USA* **95**:6169–6174

- Takahashi, N., Wang, X., Tanabe, S., Uramoto, H., Jishage, K., Uchida, S., Sasaki, S., Okada, Y. 2005. CIC-3-independent sensitivity of apoptosis to Cl⁻ channel blockers in mouse cardiomyocytes. *Cell Physiol. Biochem.* **15**:263–270
- Tallquist, M., Kazlauskas, A. 2004. PDGF signaling in cells and mice. *Cytokine Growth Factor Rev.* **15**:205–213
- Teijeiro, R., Rios, R., Costoya, J.A., Castro, R., Bello, J.L., Devesa, J., Arce, V.M. 2002. Activation of human somatostatin receptor 2 promotes apoptosis through a mechanism that is independent from induction of p53. *Cell Physiol. Biochem.* **12**:31–38
- Tohda, H., Foskett, J.K., O'Brodovich, H., Marunaka, Y. 1994. Cl⁻ regulation of a Ca(2⁺)-activated nonselective cation channel in beta-agonist-treated fetal distal lung epithelium. *Am. J. Physiol.* **266**:C104–C109
- Varela, D., Simon, F., Riveros, A., Jorgensen, F., Stutzin, A. 2004. NAD(P)H oxidase-derived H(2)O(2) signals chloride channel activation in cell volume regulation and cell proliferation. *J. Biol. Chem.* **279**:13301–13304
- Volk, T., Fromter, E., Korbmayer, C. 1995. Hypertonicity activates nonselective cation channels in mouse cortical collecting duct cells. *Proc. Natl. Acad. Sci. USA* **92**:8478–8482
- Walsh, M.F., Thamilselvan, V., Grotelueschen, R., Farhana, L., Basson, M. 2003. Absence of adhesion triggers differential FAK and SAPKp38 signals in SW620 human colon cancer cells that may inhibit adhesiveness and lead to cell death. *Cell Physiol. Biochem.* **13**:135–146
- Wang, G.L., Wang, X.R., Lin, M.J., He, H., Lan, X.J., Guan, Y.Y. 2002. Deficiency in CIC-3 chloride channels prevents rat aortic smooth muscle cell proliferation. *Circ. Res.* **91**:E28–E32
- Wang, S., Melkounian, Z., Woodfork, K.A., Cather, C., Davidson, A.G., Wonderlin, W.F., Strobl, J.S. 1998. Evidence for an early G1 ionic event necessary for cell cycle progression and survival in the MCF-7 human breast carcinoma cell line. *J. Cell Physiol.* **176**:456–464
- Wang, Z. 2004. Roles of K⁺ channels in regulating tumour cell proliferation and apoptosis. *Pfluegers Arch.* **448**:274–286
- Wehner, F., Böhmer, C., Heinzinger, H., van den, B.F., Tinel, H. 2000. The hypertonicity-induced Na(+) conductance of rat hepatocytes: physiological significance and molecular correlate. *Cell Physiol. Biochem.* **10**:335–340
- Wehner, F., Sauer, H., Kinne, R.K. 1995. Hypertonic stress increases the Na⁺ conductance of rat hepatocytes in primary culture. *J. Gen. Physiol.* **105**:507–535
- Wei, L., Xiao, A.Y., Tin, C., Yang, A., Lu, Z.Y., Yu, S.P. 2004. Effects of chloride and potassium channel blockers on apoptotic cell shrinkage and apoptosis in cortical neurons. *Pfluegers Arch.* **448**:325–334
- Wenzel, U., Daniel, H. 2004. Early and late apoptosis events in human transformed and non-transformed colonocytes are independent on intracellular acidification. *Cell Physiol. Biochem.* **14**:65–76
- Whitfield, J.F., Bird, R.P., Chakravarthy, B.R., Isaacs, R.J., Morley, P. 1995. Calcium-cell cycle regulator, differentiator, killer, chemopreventor, and maybe, tumor promoter. *J. Cell Biochem.* **22**:74–91
- Wiecha, J., Reineker, K., Reitmayer, M., Voisard, R., Hannekum, A., Mattfeldt, T., Waltenberger, J., Hombach, V. 1998. Modulation of Ca²⁺-activated K⁺ channels in human vascular cells by insulin and basic fibroblast growth factor. *Growth Horm. IGF. Res.* **8**:175–181
- Wieder, T., Essmann, F., Prokop, A., Schmelz, K., Schulze-Osthoff, K., Beyaert, R., Dorken, B., Daniel, P.T. 2001. Activation of caspase-8 in drug-induced apoptosis of B-lymphoid cells is independent of CD95/Fas receptor-ligand interaction and occurs downstream of caspase-3. *Blood* **97**:1378–1387
- Wondergem, R., Gong, W., Monen, S.H., Dooley, S.N., Gonce, J.L., Conner, T.D., Houser, M., Ecay, T.W., Ferslew, K.E. 2001. Blocking swelling-activated chloride current inhibits mouse liver cell proliferation. *J. Physiol.* **532**:661–672
- Wonderlin, W.F., Strobl, J.S. 1996. Potassium channels, proliferation and G1 progression. *J. Membrane Biol.* **154**:91–107
- Woon, L.A., Holland, J.W., Kable, E.P., Roufogalis, B.D. 1999. Ca²⁺ sensitivity of phospholipid scrambling in human red cell ghosts. *Cell Calcium* **25**:313–320
- Yu, S.P., Yeh, C.H., Sensi, S.L., Gwag, B.J., Canzoniero, L.M., Farhangrazi, Z.S., Ying, H.S., Tian, M., Dugan, L.L., Choi, D.W. 1997. Mediation of neuronal apoptosis by enhancement of outward potassium current. *Science* **278**:114–117
- Yurinskaya V.E., Goryachaya T.S., Guzhova T.V., Moshkov A.V., Rozanov Y.M., Sakuta G.A., Shirokova A.V., Shumilina E.V., Vassilieva I.O., Lang F., Vereninov A.A. 2005a. Potassium and sodium balance in U937 cells during apoptosis with and without cell shrinkage. *Cell Physiol. Biochem.* **16**: 155–162
- Yurinskaya, V.E., Moshkov, A.V., Rozanov, Y., Yu., M., Shirokova, A.V., Vassilieva, I.O., Shumilina, E.V., Lang, F., Volgareva, A.A., Vereninov, A.A. 2005b. Thymocyte K⁺, Na⁺ and water balance during dexamethasone and etoposide induced apoptosis. *Cell Physiol. Biochem.* **16**:15–22
- Zhou, Q., Kwan, H.Y., Chan, H.C., Jiang, J.L., Tam, S.C., Yao, X. 2003. Blockage of voltage-gated K⁺ channels inhibits adhesion and proliferation of hepatocarcinoma cells. *Int. J. Mol. Med.* **11**:261–266
- Zhou, Q., Zhao, J., Wiedmer, T., Sims, P.J. 2002. Normal hemostasis but defective hematopoietic response to growth factors in mice deficient in phospholipid scramblase 1. *Blood* **99**:4030–4038